




Golden Sands  
Resource Conservation  
& Development Council, Inc.



## Aquatic Invasive Species Quick Guide

### Starry Stonewort (*Nitellopsis obtusa* L.)

**Description:** Starry stonewort is a non-native species of large algae in the Characeae family. It has whorls of 4-6 long branchlets, with blunt tips. It is more robust than most members of its family, and can grow to over two meters tall. Anchored by colorless rhizoids, it contains up to several dozen 4-5mm, star-shaped bulbils. Starry stonewort typically grows in alkaline lakes with marl sediments, up to 9 meters deep. Orange reproductive structures are located in the axils of the upper branchlets. Starry stonewort is typically an annual, but can behave as a perennial during mild winters. Interestingly, starry stonewort is listed as an endangered species in the United Kingdom.



Starry stonewort has whorls of long branchlets, each with a blunt tip.

**North American Distribution:** Michigan, northern Indiana, and the northeast United States. Recently found in one lake in southeast Wisconsin.



Star-shaped bulbils give the starry stonewort its name.

**Dispersal Vectors:** Starry stonewort is native to Europe and western Asia. It was probably introduced to the Great Lakes via ballast water carried in trans-oceanic ships. Fragments of starry stonewort can easily be spread between lakes by boats, trailers, and anchors holding sediments. Local dispersal occurs by bulbils being transported by water currents or animals within the lake.

**Ecological Impacts:** By forming extremely dense mats of vegetation, starry stonewort can greatly reduce the diversity of aquatic plants in a lake. It can also impede movement of fish and other animals, and can decrease successful spawning activity. Mats growing to the surface can reduce water flow and prevent recreational activities.

**Control Options:** Manual removal of starry stonewort is difficult and probably impractical on a large scale. Abundant bulbils on the rhizoids can dislodge if disturbed, and will sprout new individuals. Manual removal efforts must emphasize careful removal of these bulbils.

Copper-based herbicides have been effective at suppressing starry stonewort. Endothall is sometimes added to copper herbicides to increase its effectiveness. Herbicide applications may be less effective on tall stands of starry stonewort, as the chemical is quickly absorbed into the upper parts of the algae, leaving the lower parts unharmed. Most states require chemical use permits for any herbicide treatments in standing water or wetland situations.

An effective biological control agent is not known at this time.

#### Additional Information:

Pullman, G. Douglas and Gary Crawford. 2010. A decade of starry stonewort in Michigan. *Lakeline*. 36-42.

**Photo credit:** Paul Skawinski



Starry stonewort (front, center), much more robust than the surrounding native muskgrasses (*Chara* spp.).

This Quick Guide is part of a series on aquatic invasive species, and may be reproduced for educational purposes. Visit us at [www.uwsp.edu/uwexlakes/clmn](http://www.uwsp.edu/uwexlakes/clmn) or [www.goldensandsrscd.org/our-work/water](http://www.goldensandsrscd.org/our-work/water) to download this series of handouts.

Developed by Golden Sands Resource Conservation & Development (RC&D) Council, Inc. as part of an aquatic invasive species (AIS) education program, supported by an AIS grant from the Wisconsin Department of Natural Resources.

## Starry stonewort *Nitellopsis obtusa*

### Description

This species is listed on the Michigan prohibited aquatic plant list. Appears in July and increases through October.

### Habit

Aquatic macroalga with smooth, irregular branches; rootlike rhizoids. Forms dense mats that cover lake bottoms.

### Leaves

Whorls of 5-8 fine, uneven-length branches arise at an acute angle from nodes. Contents can be squeezed out of tube-like cell wall when cut; dark reddish gametangia.

### Stems

May reach 80 cm (33 in) in length.

### Flowers



Lacks true flowers. Cream colored bulbs form at the base of each cluster of branches.

### **Fruits and Seeds**

### **Habitat**

Native to Europe and Asia. Found in still or slow moving waters; grows at depths ranging from 1 m (39 in) to 30 m (~93 ft).

### **Reproduction**

Probably introduced in ballast water. Dispersed by boats, waterfowl.

### **Similar**

Common stonewort (*Chara vulgaris*), which feels brittle and scaly, unlike the starry stonewort.

### **Monitoring and Rapid Response**

Algaecides and mechanical harvesting.

### **Credits**

The information provided in this factsheet was gathered from the Michigan Lake Info and the Michigan Natural Features Inventory. Individual species images that appear with a number in a black box are courtesy of the Bugwood.org network (<http://www.invasive.org>). Individual photo author credits are not included due to the small display size of the images and subsequent difficulty of reading the provided text. All other images appear courtesy of Google (<http://images.google.com>).